

América do Norte

Antrozous pallidus (Pallid), Corynorhinus townsendii (Townsend's big-eared bat), Eptesicus fuscus (Big brown), Euderma maculatum (Spotted), Eumops floridanus (Bonneted), Eumops perotis (Western mastiff), Eumops underwoodi (Underwood's bonneted), Lasionycteris noctivagans (Silver-haired), Lasiurus blossevilli (Western red), Lasiurus borealis (Eastern red), Lasiurus cinereus (Hoary), Lasiurus ega (Southern yellow), Lasiurus intermedius (Northern yellow), Lasiurus seminolus (Seminole bat), Lasiurus xanthinus (Western yellow), Macrotus californicus (California leaf-nosed), Molossus molossus (Pallas's mastiff), Mormoops megalophylla (Ghost faced), Myotis austroriparius (Southeastern myotis), Myotis californicus (California myotis), Myotis ciliolabrum (Western Small-footed), Myotis evotis (Long-eared myotis), Myotis grisescens (Gray), Myotis leibii (Small-footed), Myotis lucifugus (Little brown), Myotis occultus (Arizona myotis), Myotis septentrionalis (Northern long-eared myotis), Myotis sodalis (Indiana), Myotis thysanodes (Fringed myotis), Myotis velifer (Cave myotis), Myotis volans (Long-legged myotis), Myotis yumanensis (Yuma myotis), Nycticeius humeralis (Evening), Nyctinomops femorosaccus (Pocketed free-tailed), Nyctinomops macrotis (Big free-tailed), Parastrellus hesperus (Western pipistrelle), Perimyotis subflavus (Tricolored), Tadarida brasiliensis (Mexican free-tailed)

Reino unido e Europa

Barbastella barbastellus (Western barbastelle)*, Eptesicus isabellinus (Meridional serotine), Eptesicus isabellinus (Meridional serotine), Eptesicus nilssonii (Northern), Eptesicus serotinus (Serotine)*, Hypsugo savii (Savi's pipistrelle), Miniopterus schreibersii (Common bent-wing), Miniopterus schreibersii (Common bent-wing), Myotis alcathoe (Alcathoe's), Myotis bechsteinii (Bechstein's)*, Myotis brandtii (Brandt's)*, Myotis capaccinii (Long-fingered), Myotis capaccinii (Long-fingered), Myotis dasycneme (Pond), Myotis daubentonii (Daubenton's)*, Myotis emarginatus (Geoffroy's), Myotis escaleraei (Escalera's), Myotis escaleraei (Escalera's), Myotis myotis (Greater mouse-eared), Myotis mystacinus (Whiskered), Myotis nattereri (Natterer's)*, Nyctalus lasiopterus (Greater noctule), Nyctalus leisleri (Lesser noctule)*, Nyctalus noctula (Common noctule)*, Pipistrellus kuhlii (Kuhl's pipistrelle), Pipistrellus nathusii (Nathusius's pipistrelle)*, Pipistrellus pipistrellus (Common pipistrelle)*, Pipistrellus pygmaeus (Soprano pipistrelle)*, Plecotus auritus (Brown long eared)*, Plecotus austriacus (Grey long-eared), Plecotus austriacus (Grey long-eared), Rhinolophus euryale (Mediterranean horseshoe), Rhinolophus ferrumequinum (Greater horseshoe)*, Rhinolophus hipposideros (Lesser horseshoe)*, Tadarida teniotis (European free-tailed), Tadarida teniotis (European free-tailed), Vespertilio murinus (Particoloured)

Neotrópicos

Antrozous pallidus (Pallid), Balantiopteryx io (Thomas's sac-winged), Balantiopteryx plicata (Gray sac-winged), Balantiopteryx plicata (Gray sac-winged), Bauerus dubiaquercus (Van Gelder's), Centronycteris centralis (Thomas's shaggy), Centronycteris maximiliani (Shaggy), Corynorhinus townsendii (Townsend's big-eared), Cynomops mexicanus (Mexican dog-faced), Diclidurus albus (Northern ghost), Eptesicus brasiliensis (Brazilian brown), Eptesicus furinalis (Argentine brown), Eptesicus fuscus (Big brown), Euderma maculatum (Spotted), Eumops glaucinus (Eumops floridanus), Eumops perotis (Western Mastiff), Eumops underwoodi (Underwood's bonneted), Lasionycteris noctivagans (Silver haired), Lasionycteris noctivagans (Silver-haired), Lasiurus blossevillii (Desert red), Lasiurus borealis (Eastern red), Lasiurus cinereus (Hoary), Lasiurus ega (Southern yellow), Lasiurus insularis (Cuban Yellow), Lasiurus intermedius (Northern yellow), Lasiurus seminolus (Seminole), Lasiurus xanthinus (Western yellow), Lasiurus xanthinus (Western yellow), Macrotus californicus (California leaf-nosed), Macrotus californicus (California leaf-nosed), Molossus molossus (Velvety free-tailed), Molossus rufus (Black mastiff), Molossus sinaloae (Sinaloan mastiff), Myotis occultus (Arizona myotis), Myotis velifer (Cave myotis), Nyctinomops femorosaccus (Pocketed free-tail), Nyctinomops macrotis (Big free-tailed)

África do Sul

Chaerophon pumilus (Little free-tailed), Eptesicus hottentotus (Long-tailed house), Laephotis botswanae (Botswanan long-eared), Miopterus natalensis (Natal long-fingered), Myotis bocagei (Rufous mouse-eared), Neoromicia capensis (Cape serotine), Pipistrellus hesperidus (Dusky pipistrelle), Rhinolophus blasii (Blasius's horseshoe), Rhinolophus capensis (Cape horseshoe), Rhinolophus clivosis (Geoffroy's horseshoe), Rhinolophus denti (Dent's horseshoe), Rhinolophus fumigatus (Rüppell's horseshoe), Rhinolophus hildebrandtii (Hildebrandt's horseshoe), Rhinolophus landeri (Lander's horseshoe), Rhinolophus simulator (Bushveld horseshoe), Rhinolophus smithersi (Smithers's horseshoe), Rhinolophus swinnyi (Swinny's horseshoe), Sauromys petrophilus (Roberts's flat-headed), Scotophilus dinganii (African yellow), Scotophilus Nigrita (Schreber's yellow), Tadarida aegyptiaca (Egyptian free-tailed)

NOTA: O Auto ID destina-se ao uso na análise de gravações de morcegos isolados em vôo livre em ambientes com pouca interferência. Gravações em aglomerados, múltiplos indivíduos, morcegos em ambientes com alta interferência ou chamadas sociais não são adequados para identificação automática e os resultados podem não ser precisos.